

5. (THEORY OF) GW DETECTION

GW predicted in 1915ish. But researchers were not sure they were real until the 1960s!
 → in GR we cannot use coordinates to define observers, so can GW be transformed away with a coordinate transformation

FEYNMAN "STICKY BEAD" ARGUMENT (CHAPEL HILL CONFERENCE 1957)

Feynman's gravitational wave detector: It is simply two beads sliding freely (but with a small amount of friction) on a rigid rod. As the wave passes over the rod, atomic forces hold the length of the rod fixed, but the proper distance between the two beads oscillates. Thus, the beads rub against the rod, dissipating heat. -Jin

Amazing book on the history of GW:

TRAVELING AT THE SPEED OF THOUGHT by KENNEFICK

In current GW research: what is the eccentricity of a binary? We cannot use the shape of the orbit, that's gauge dependent!
 BOSCHINI, LAUREL, GEROSA, FUMAGALLI (2025)
 SHAIKH, VARMA et al (2023)

Interaction of a GW with test masses
 ↳ eg the LIGO mirror!

GEODESIC EQUATION IN GR $ds^2 = g_{\mu\nu} dx^\mu dx^\nu$ SEC 1.3

$$\sim \frac{dx^\mu}{d\tau^2} + \Gamma_{\nu\rho}^\mu(x) \frac{dx^\nu}{d\tau} \frac{dx^\rho}{d\tau} = 0 \quad (1.68)$$

GEODESIC DEVIATION Two geodesics $x^\mu(\tau)$, $x^\mu(\tau) + \xi^\mu(\tau)$

$$\sim \frac{d^2 \xi^\mu}{d\tau^2} + 2\Gamma_{\nu\rho}^\mu(x) \frac{dx^\nu}{d\tau} \frac{d\xi^\rho}{d\tau} + \xi^\sigma \partial_\sigma \Gamma_{\nu\rho}^\mu(x) \frac{dx^\nu}{d\tau} \frac{dx^\rho}{d\tau} = 0 \quad (1.69)$$

TT FRAME ... A somewhat special frame where GW have simple expression ... what's physically?

Take a test mass at rest at $\tau = 0$

$$\sim \frac{d^2 x^i}{d\tau^2} \Big|_{\tau=0} = - \left[\Gamma_{\infty}^i \left(\frac{dx^0}{d\tau} \right)^2 \right]_{\tau=0} \quad (1.77)$$

in linearized gravity $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$

$$P_i \approx \frac{1}{2} (2\dot{x}_0 h_{0i} - \dot{x}_i h_{00})$$

but in the TT gauge $h_{0i} = h_{00} = 0 \Rightarrow \left. \frac{dx^i}{dt} \right|_{t=0} = 0$

\Rightarrow in the TT gauge $\frac{dx^i}{dt} = 0$ and stays zero
particles remain at rest before/during/after
the arrival of GW

The TT gauge is the frame where coordinates
stretch and squeeze appropriately such that
they do not change

Implementation: freely falling test masses where
the masses themselves are used to mark the
coordinates

DETECTOR FRAME

In a lab positions are not marked by freely falling
test masses. One ideally uses a "rigid ruler"
to mark coordinates

\rightarrow we do not measure things in the TT frame!

Drop free satellite in free fall. Make measurements
in test lab.

Take a sufficiently small region of size x^i

Local inertial frame:

$$ds^2 \approx -dt^2 + \delta_{ij} dx^i dx^j$$

Expansion of $g_{\mu\nu} dx^\mu dx^\nu$ to 2nd order in x^i
 $\approx ds^2 \approx -dt^2 (1 + R_{0i0j} x^i x^j) - 2dt dx^i \left(\frac{2}{3} R_{0jik} x^j x^k \right)$
 $+ dx^i dx^j \left(\delta_{ij} - \frac{1}{3} R_{ijkl} x^k x^l \right)$ (1.57)

For a detector on Earth (say LIGO...) we have an acceleration $\bar{a} = -\bar{g}$ and a rotation $\bar{\Omega}$ (Foucault's pendulum...)

$$ds^2 \approx -dt^2 \left[1 + 2\bar{a}\bar{x} + (\bar{a}\bar{x})^2 - (\bar{\Omega} \times \bar{x}) + R_{0ij} x^i x^j \right] + 2dt dx^i \left[\epsilon_{ijk} \bar{\Omega}^j x^k - \frac{2}{3} R_{0ijk} x^j x^k \right] + dx^i dx^j \left[\delta_{ij} - \frac{1}{3} R_{ikje} x^k x^e \right] \quad (1.88)$$

↳ this is called "proper lab frame" and it's where we make measurements.

$2\bar{a}\bar{x} \rightarrow$ inertial acceleration

$(\bar{a}\bar{x})^2 \rightarrow$ gravitational redshift

$(\bar{\Omega} \times \bar{x}) \rightarrow$ Lorentz time dilation

$\epsilon_{ijk} \bar{\Omega}^j x^k \rightarrow$ Sagnac effect

$R_{ikje} \rightarrow$ both slowly varying perturbation field and the GWs

Given this metric, the geodesic equation is

$$\frac{d^2 x^i}{dt^2} = -a^i - 2(\bar{\Omega} \times \bar{v})^i + \frac{F^i}{m} + o(x^i) \quad (1.89)$$

gravity

Coriolis

extended forces (say the LIGO suspension system)

\rightarrow centrifugal acceleration $\bar{\Omega} \times (\bar{\Omega} \times \bar{r})$ is of $o(x^i)$

\rightarrow GWs are in the $o(x^i)$ term as well

m. particle find a frequency region where the slowly varying gravitational field is not important, sufficient isolation, for LIGO, seismic between 10Hz and 100Hz
 \approx let's focus on this frequency range,

only the $G_{\mu\nu}$ are present in the R terms
 Geodesic deviation gives

$$\frac{d^2 \xi^i}{d\tau^2} + \xi^\sigma \partial_\sigma R^i_{0j0} \left(\frac{dx^0}{d\tau}\right)^2 = 0 \quad (1.9p) \quad (\text{at the expansion point } \dot{r}^\sigma = 0)$$

$$\frac{d^2 \xi^i}{d\tau^2} = -R^i_{0j0} \xi^j \left(\frac{dx^0}{d\tau}\right)^2$$

Ass at rest initially, then velocity $\frac{dx^i}{d\tau} = 0(a)$

$$R \sim 0(a)$$

$$d\tau \approx dt (1 + 0(a))$$

so we can write to linear order

$$\frac{d^2 \xi^i}{dt^2} = -R^i_{0j0} \xi^j$$

R tensor is invariant (not just covariant) in linearized gravity (proof around (4.13) in MASSIVE)
 so I can compute it in the easy π gauge

$$R^i_{0j0} = -\frac{1}{2} \ddot{h}^{\pi}_{ij}$$

$$\Rightarrow \ddot{\xi}^i = \frac{1}{2} \ddot{h}^{\pi}_{ij} \xi^j$$

this is what we actually measure!

That is, a Newtonian description with a force acting on the mirror

$$F_i = \frac{m}{2} \ddot{h}^{\pi}_{ij} \xi^j$$

- Note ξ and t are coordinate separation (which we measure), not proper distance
- only in the direction where the mirror is free to move

We have expanded everything to a "small" region where our experiment beaver. OK only if
 size $L \ll \lambda \rightarrow$ GW wavelength

OK for LIGO, not for LISA


(admittedly, I don't understand the Fabry-Pérot cavity business in this...)

DETECTOR RESPONSE

compute the response of an interferometer with 90° opening angle and equal arms, like LIGO, VIRGO, KAGRA

$$\ddot{\xi}^i = \frac{1}{2} \ddot{h}_{ij} \xi^j$$

\rightarrow Mirror located at $(L, 0, 0)$
 we're interested in the motion along x only $\ddot{\xi}_x = \frac{1}{2} \ddot{h}_{xx} L$




\rightarrow Mirror located at $(0, L, 0)$, motion only along y $\ddot{\xi}_y = \frac{1}{2} \ddot{h}_{yy} L$

Say the GW comes from the z axis, orthogonal to the detector. We have $h_{xx} = h_+$ $h_{yy} = -h_+$ (see Secter 1)

$$\Rightarrow h_+ = \frac{1}{2} (h_{xx} - h_{yy})$$

In general, for a generic direction we must compute $\frac{1}{2} (h_{xx} - h_{yy}) \dots$

Direction of propagation z' that forms a polar angle θ with z and an azimuthal angle ψ in the plane of the detector



$$a'_{ij} = \begin{pmatrix} a_+ & a_x & 0 \\ a_x & -a_+ & 0 \\ 0 & 0 & 0 \end{pmatrix}_{ij}$$

$$a_{ij} = R_z(\phi) R_y(\theta) a'_{ij} \quad \text{rotation matrices}$$

$$\Rightarrow a_{xx} = a_+ (\cos^2 \theta \cos^2 \phi - \sin^2 \phi) + 2 a_x \cos \theta \sin \phi \cos \phi \quad (9.133)$$

$$a_{yy} = a_+ (\cos^2 \theta \sin^2 \phi - \cos^2 \phi) - 2 a_x \cos \theta \sin \phi \cos \phi \quad (9.144)$$

$$a(t) = \frac{1}{2} (a_{xx} - a_{yy}) = \frac{1}{2} a_+ (1 + \cos^2 \theta) \cos 2\phi + a_x \cos \theta \sin 2\phi$$

only for an interferometer
we will be response of a detector simply $a(t)$...

The above a'_{ij} shows that a_+ is along x' . But an extra degree of freedom ψ , which is a rotation in the $x'y'$ plane. More rotation matrices... (see PASSIVE problem 2.1). The result is

$$a(t) = F_+(\theta, \phi, \psi) a_+ + F_x(\theta, \phi, \psi) a_x$$


BEAM PATTERN OF A QW INTERFEROMETER

$$F_+ = \frac{1}{2} (1 + \cos^2 \theta) \cos 2\phi \cos 2\psi - \cos \theta \sin 2\phi \sin 2\psi$$

$$F_x = \frac{1}{2} (1 + \cos^2 \theta) \cos 2\phi \sin 2\psi + \cos \theta \sin 2\phi \cos 2\psi$$

θ, ϕ : sky location of the source

ψ : polarization angle



+ polarization

x polarization

- Interferometer is maximally sensitive for a source aligned
- Interferometer is blind for a source along the arm.

Caution: do not get confused with the emission pattern of a binary...

SEE APPENDIX 1
OF GEOSA PAPER
VECHIO 2020

The binary waveform is

$$a_+(t) = A(t) \frac{1 + \cos^2 i}{2} \cos \phi(t)$$

$$a_x(t) = A(t) \cos i \sin \phi(t)$$

$A(t)$: GW amplitude

$\phi(t)$: GW phase

i : inclination

\leadsto we can write (FINN CHERNOFF 1993)

$$a(t) = \omega A(t) \cos[\phi(t) - \phi_0]$$

$$\text{with } \omega = \sqrt{\left(F_+ \frac{1 + \cos^2 i}{2}\right)^2 + (F_x \cos i)^2}$$

PROJECTION
FACTOR

$$\phi_0 = \arctan \frac{2F_x \cos i}{F_+ (1 + \cos^2 i)}$$

PHASE SHIFT
(independent of constant
measures \mathcal{L})

The various angles are all into ω

One has $0 < \omega < 1$ and $\omega = 1$ only for:

$\theta = 0$: source overhead

$i = 0$: source face-on